

Preprint, arXiv:1010.4298

FAST CONVERGING SERIES FOR SOME FAMOUS CONSTANTS

ZHI-WEI SUN

Department of Mathematics, Nanjing University
Nanjing 210093, People's Republic of China
zwsun@nju.edu.cn
<http://math.nju.edu.cn/~zwsun>

ABSTRACT. For $k = 0, 1, 2, \dots$ let $H_k^{(2)} := \sum_{0 < j \leq k} 1/j^2$ and $\bar{H}_k^{(2)} := \sum_{0 < j \leq k} 1/(2j-1)^2$. In this paper we give a new type of exponentially converging series (involving $H_k^{(2)}$ or $\bar{H}_k^{(2)}$) for certain famous constants. For example, we show that

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{H_{k-1}^{(2)}}{k^2 \binom{2k}{k}} &= \frac{\pi^4}{1944}, \quad \sum_{k=1}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)}}{(2k+1)(-16)^k} = -\frac{1}{3} \log^3 \phi, \\ \sum_{k=1}^{\infty} \frac{L_{2k} H_{k-1}^{(2)}}{k^2 \binom{2k}{k}} &= \frac{41\pi^4}{7500}, \quad \sum_{k=1}^{\infty} \frac{\binom{2k}{k} L_{2k+1} \bar{H}_k^{(2)}}{(2k+1)16^k} = \frac{13\pi^3}{1500}, \end{aligned}$$

where ϕ is the golden ratio $(\sqrt{5}+1)/2$, and L_0, L_1, L_2, \dots are Lucas numbers given by $L_0 = 2$, $L_1 = 1$, and $L_{n+1} = L_n + L_{n-1}$ ($n = 1, 2, 3, \dots$).

1. INTRODUCTION

Series for powers of π or $\log 2$ have a long history (see, e.g., [M]). It is well known that

$$\sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} = \frac{\pi}{4}, \quad \zeta(2) := \sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}, \quad \zeta(4) := \sum_{k=1}^{\infty} \frac{1}{k^4} = \frac{\pi^4}{90}$$

and

$$\log 2 = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k}.$$

2010 *Mathematics Subject Classification*. Primary 33B99, 65B10; Secondary 05A10, 05A19, 11A07, 11B39, 11B65, 11Y60.

Keywords. Series for powers of π , central binomial coefficients, Fibonacci and Lucas numbers, golden ratio, congruences.

Supported by the National Natural Science Foundation (grant 10871087) and the Overseas Cooperation Fund (grant 10928101) of China.

However, these series converge very slow. In 1974 Gosper announced the following fast converging series for π (cf. G. Almkvist, C. Krattenthaler and J. Petersson [AKP]):

$$\sum_{k=0}^{\infty} \frac{25k-3}{2^k \binom{3k}{k}} = \frac{\pi}{2}.$$

(Note that $n! \sim \sqrt{2\pi n}(n/e)^n$ ($n \rightarrow +\infty$) by Stirling's formula.) In 1993 D. Zeilberger [Z] obtained the identity

$$\sum_{k=1}^{\infty} \frac{21k-8}{k^3 \binom{2k}{k}^3} = \frac{\pi^2}{6}$$

via the WZ method (see, e.g., [PWZ]). This is a rapidly converging series for π^2 since

$$\binom{2k}{k} \sim \frac{4^k}{\sqrt{k\pi}} \quad (k \rightarrow +\infty)$$

by Stirling's formula. In 2006 D. Bailey, J. Borwein and D. Bradley [BBB] obtained the celebrated formula

$$\sum_{k=1}^{\infty} \frac{1}{k^4 \binom{2k}{k}} = \frac{17}{36} \zeta(4) = \frac{17\pi^4}{3240}.$$

Here is a fast converging series for $\log 2$ given by Borwein and Bailey [BB. p. 129]:

$$\log 2 = \frac{2}{3} \sum_{k=0}^{\infty} \frac{1}{(2k+1)9^k}.$$

Recall that harmonic numbers are those rational numbers

$$H_n := \sum_{0 < k \leq n} \frac{1}{k} \quad (n = 0, 1, 2, \dots).$$

and *harmonic numbers of the second order* are defined by

$$H_n^{(2)} := \sum_{0 < k \leq n} \frac{1}{k^2} \quad (n = 0, 1, 2, \dots).$$

Recently the author [S6] found and proved that

$$\sum_{k=1}^{\infty} \frac{2^k H_{k-1}^{(2)}}{k \binom{2k}{k}} = \frac{\pi^3}{48},$$

which was motivated by his philosophy of correspondence between certain series and congruences (cf. [S5]) and his related result

$$\sum_{k=1}^{p-1} \frac{\binom{2k}{k}}{2^k} H_k^{(2)} \equiv -E_{p-3} \pmod{p} \quad \text{for any prime } p > 3,$$

where E_0, E_1, E_2, \dots are Euler numbers given by $E_0 = 1$ and the recursion

$$\sum_{\substack{k=0 \\ 2|k}}^n \binom{n}{k} E_{n-k} = 0 \quad (n = 1, 2, 3, \dots).$$

The reader may consult [PS], [S1-6] and [ST1, ST2] for congruences involving central binomial coefficients which provide important backgrounds for our current research on series. In [S6] the author explained how he was led to consider $\sum_{k=0}^{p-1} \binom{2k}{k} H_k^{(2)} / 2^k$ modulo an odd prime p and the evaluation of the series $\sum_{k=1}^{\infty} 2^k H_{k-1}^{(2)} / (k \binom{2k}{k})$. Note that $H_n^{(2)} \rightarrow \zeta(2) = \pi^2/6$ as $n \rightarrow +\infty$.

In this paper we aim at giving certain exponentially converging series for some classical constants (such as π^3 and π^4) with each term related to the second-order harmonic numbers.

Recall that the Fibonacci sequence $\{F_n\}_{n \geq 0}$ and the Lucas sequence $\{L_n\}_{n \geq 0}$ are given by

$$F_0 = 0, F_1 = 1, \text{ and } F_{n+1} = F_n + F_{n-1} \quad (n = 1, 2, 3, \dots),$$

and

$$L_0 = 2, L_1 = 1, \text{ and } L_{n+1} = L_n + L_{n-1} \quad (n = 1, 2, 3, \dots).$$

It is well known that for each $n = 0, 1, 2, \dots$ we have

$$\sqrt{5}F_n = \left(\frac{1+\sqrt{5}}{2}\right)^n - \left(\frac{1-\sqrt{5}}{2}\right)^n \quad \text{and} \quad L_n = \left(\frac{1+\sqrt{5}}{2}\right)^n + \left(\frac{1-\sqrt{5}}{2}\right)^n.$$

Now we state our first theorem in which we relate Fibonacci numbers and Lucas numbers to π^4 for the first time.

Theorem 1.1. (i) *We have the new identity*

$$\sum_{k=1}^{\infty} \frac{H_{k-1}^{(2)}}{k^2 \binom{2k}{k}} = \frac{\pi^4}{1944}. \quad (1.1)$$

Also,

$$\sum_{k=1}^{\infty} \frac{L_{2k} H_{k-1}^{(2)}}{k^2 \binom{2k}{k}} = \frac{41\pi^4}{7500}, \quad \sum_{k=1}^{\infty} \frac{F_{2k} H_{k-1}^{(2)}}{k^2 \binom{2k}{k}} = \frac{2\pi^4}{375\sqrt{5}}, \quad (1.2)$$

and

$$\sum_{k=1}^{\infty} \frac{v_k H_{k-1}^{(2)}}{k^2 \binom{2k}{k}} = \frac{34\pi^4}{1875} \quad \text{and} \quad \sum_{k=1}^{\infty} \frac{u_k H_{k-1}^{(2)}}{k^2 \binom{2k}{k}} = \frac{2\pi^4}{125\sqrt{5}}, \quad (1.3)$$

where

$$u_k = \begin{cases} 5^{k/2} F_k & \text{if } 2 \mid k, \\ 5^{(k-1)/2} L_k & \text{if } 2 \nmid k, \end{cases} \quad \text{and} \quad v_k = \begin{cases} 5^{k/2} L_k & \text{if } 2 \mid k, \\ 5^{(k+1)/2} F_k & \text{if } 2 \nmid k. \end{cases}$$

(ii) We also have

$$\sum_{k=1}^{\infty} \frac{H_{k-1}^{(2)} x^k}{k \binom{2k}{k}} = \frac{4}{3} \sqrt{\frac{x}{4-x}} \arcsin^3 \frac{\sqrt{x}}{2} \quad \text{for } 0 \leq x < 4. \quad (1.4)$$

In particular,

$$\sum_{k=1}^{\infty} \frac{H_{k-1}^{(2)}}{k \binom{2k}{k}} = \frac{\pi^3}{162\sqrt{3}} \quad (1.5)$$

and

$$\sum_{k=1}^{\infty} \frac{3^k H_{k-1}^{(2)}}{k \binom{2k}{k}} = \frac{4\pi^3}{27\sqrt{3}}. \quad (1.6)$$

Remark 1.1. (a) The modern **Mathematica 7** (version 7) employs various powerful algorithms (including the WZ method) to obtain summations of series, however it is still unable to evaluate the series in Theorem 1.1. (b) Motivated by massive Feynman diagrams in physics, A. I. Davydychev and M. Yu. Kalmykov [DK, (C16) and (C17)] obtained that for any $x > 1/4$ we have

$$\sum_{k=1}^{\infty} \frac{H_{k-1}^{(2)}}{k \binom{2k}{k} (-x)^k} = \frac{1}{6\sqrt{4x+1}} \log^3 \frac{\sqrt{4x+1}+1}{\sqrt{4x+1}-1}$$

and

$$\sum_{k=1}^{\infty} \frac{H_{k-1}^{(2)}}{k^2 \binom{2k}{k} (-x)^k} = \frac{1}{24} \log^4 \frac{\sqrt{4x+1}+1}{\sqrt{4x+1}-1}.$$

For convenience, we set

$$\bar{H}_n^{(2)} := \sum_{0 < k \leq n} \frac{1}{(2k-1)^2} = H_{2n}^{(2)} - \frac{H_n^{(2)}}{4} \quad \text{for } n = 0, 1, 2, \dots$$

Our second theorem and its corollaries deal with exponentially converging series for π^3 , π^2 or $\log^3 x$ with the k th term related to $\bar{H}_k^{(2)}$.

Theorem 1.2. (i) If $0 < x \leq 4$, then

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)}}{2k+1} \left(\frac{x}{16}\right)^k = \frac{\arcsin^3(\sqrt{x}/2)}{3\sqrt{x}}. \quad (1.7)$$

Also,

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k} F_{2k+1} \bar{H}_k^{(2)}}{(2k+1)16^k} = \frac{7\pi^3}{750\sqrt{5}} \quad (1.8)$$

and

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k} L_{2k+1} \bar{H}_k^{(2)}}{(2k+1)16^k} = \frac{13\pi^3}{1500}. \quad (1.9)$$

(ii) For $x \geq 1$ we have

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)}}{(2k+1)(-4x)^k} = \frac{\sqrt{x}}{6} \log^3 \frac{\sqrt{x+1}-1}{\sqrt{x}}. \quad (1.10)$$

In view of (1.7) and (1.10), one can easily deduce the following consequence.

Corollary 1.1. We have

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)}}{(2k+1)16^k} = \frac{\pi^3}{648}, \quad (1.11)$$

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)}}{(2k+1)8^k} = \frac{\pi^3}{192\sqrt{2}}, \quad (1.12)$$

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)}}{2k+1} \left(\frac{3}{16}\right)^k = \frac{\pi^3}{81\sqrt{3}}, \quad (1.13)$$

and

$$\sum_{k=0}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)}}{(2k+1)(-32)^k} = -\frac{\sqrt{2}}{24} \log^3 2, \quad (1.14)$$

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)}}{(2k+1)(-12)^k} = -\frac{\sqrt{3}}{48} \log^3 3, \quad (1.15)$$

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)}}{(2k+1)(-16)^k} = -\frac{1}{3} \log^3 \frac{\sqrt{5}+1}{2}. \quad (1.16)$$

Remark 1.2. (1.11) also follows from Section 2 of [PP]. Our (1.16) provides a fast converging series for $\log^3 \phi$ (at a geometric rate with ratio $-1/4$), where $\phi = (\sqrt{5}+1)/2$ is the famous golden ratio. It seems that there are no previously known exponentially converging series for $\log^3 \phi$.

Corollary 1.2. *For $0 \leq x < 2$ we have*

$$\sum_{k=1}^{\infty} \binom{2k}{k} \bar{H}_k^{(2)} \left(\frac{x}{4}\right)^{2k} = \frac{\arcsin^2(x/2)}{\sqrt{4-x^2}}. \quad (1.17)$$

In particular,

$$\sum_{k=1}^{\infty} \binom{2k}{k} \frac{\bar{H}_k^{(2)}}{16^k} = \frac{\pi^2}{36\sqrt{3}}, \quad (1.18)$$

$$\sum_{k=1}^{\infty} \binom{2k}{k} \frac{\bar{H}_k^{(2)}}{8^k} = \frac{\pi^2}{16\sqrt{2}}, \quad (1.19)$$

$$\sum_{k=1}^{\infty} \binom{2k}{k} \frac{3^k \bar{H}_k^{(2)}}{16^k} = \frac{\pi^2}{9}. \quad (1.20)$$

To conclude this section, we raise a conjecture on related congruences.

Conjecture 1.1. *Let p be an odd prime.*

(i) *If $p > 3$ then*

$$\sum_{k=1}^{p-1} \frac{\binom{2k}{k}}{k} H_k^{(2)} \equiv \frac{2H_{p-1}}{3p^2} + \frac{76}{135} p^2 B_{p-5} \pmod{p^3},$$

where B_0, B_1, B_2, \dots are well known Bernoulli numbers. Also,

$$\sum_{k=1}^{p-1} \frac{\binom{2k}{k}}{8^k} \bar{H}_k^{(2)} \equiv \left(\frac{-2}{p}\right) \frac{E_{p-3}}{4} \pmod{p},$$

where $(-)$ denotes the Legendre symbol.

(ii) *If $p > 3$, then*

$$\sum_{k=1}^{p-1} (-2)^k \binom{2k}{k} H_k^{(2)} \equiv \frac{2}{3} q_p(2)^2 \pmod{p},$$

where $q_p(2)$ denotes the Fermat quotient $(2^{p-1} - 1)/p$. When $p > 5$, we have

$$\sum_{k=1}^{p-1} (-1)^k \binom{2k}{k} H_k^{(2)} \equiv \frac{5}{2} \left(\frac{p}{5}\right) \frac{F_{p-(\frac{p}{5})}^2}{p^2} \pmod{p}.$$

2. PROOFS OF THEOREMS 1.1-1.2 AND COROLLARY 1.2

Proof of Theorem 1.1. By **Mathematica 7**, for $0 \leq x \leq 4$ we have

$$\sum_{k=1}^{\infty} \frac{H_{k-1}^{(2)} x^k}{k^2 \binom{2k}{k}} = \frac{2}{3} \arcsin^4 \frac{\sqrt{x}}{2}. \quad (2.1)$$

Putting $x = 1$ in (2.1) we get (1.1), however the series cannot be evaluated by **Mathematica 7** via the **FullSimplify** command.

Let $\alpha = (1 + \sqrt{5})/2$ and $\beta = (1 - \sqrt{5})/2$. Then

$$\arcsin \frac{\sqrt{\alpha^2}}{2} = \arcsin \frac{\sqrt{5} + 1}{4} = \frac{3\pi}{10}$$

and

$$\arcsin \frac{\sqrt{\beta^2}}{2} = \arcsin \frac{\sqrt{5} - 1}{4} = \frac{\pi}{10}.$$

Therefore

$$\sum_{k=1}^{\infty} \frac{H_{k-1}^{(2)} \alpha^{2k}}{k^2 \binom{2k}{k}} = \frac{2}{3} \left(\frac{3\pi}{10} \right)^4 \quad \text{and} \quad \sum_{k=1}^{\infty} \frac{H_{k-1}^{(2)} \beta^{2k}}{k^2 \binom{2k}{k}} = \frac{2}{3} \left(\frac{\pi}{10} \right)^4.$$

As $\alpha^{2k} + \beta^{2k} = L_{2k}$ and $\alpha^{2k} - \beta^{2k} = \sqrt{5} F_{2k}$, (1.2) follows.

It is easy to see that for any $k = 0, 1, 2, \dots$ we have

$$\sqrt{5} u_k = \left(\frac{5 + \sqrt{5}}{2} \right)^k - \left(\frac{5 - \sqrt{5}}{2} \right)^k \quad \text{and} \quad v_k = \left(\frac{5 + \sqrt{5}}{2} \right)^k + \left(\frac{5 - \sqrt{5}}{2} \right)^k.$$

Also,

$$\sin \frac{2\pi}{5} = \sqrt{1 - \left(\frac{\sqrt{5} - 1}{4} \right)^2} = \frac{1}{2} \sqrt{\frac{5 + \sqrt{5}}{2}}$$

and

$$\sin \frac{\pi}{5} = \sqrt{1 - \left(\frac{\sqrt{5} + 1}{4} \right)^2} = \frac{1}{2} \sqrt{\frac{5 - \sqrt{5}}{2}}.$$

Thus, applying (2.1) with $x = (5 \pm \sqrt{5})/2$ we obtain (1.3) immediately.

Since $\binom{2k}{k} \sim 4^k / \sqrt{k\pi}$ as $k \rightarrow \infty$, the series $\sum_{k=1}^{\infty} 4^k / (k^2 \binom{2k}{k})$ converges.

Note also that $H_{k-1}^{(2)} < \zeta(2) = \pi^2/6$ for $k = 1, 2, 3, \dots$. Thus, we can take derivatives of both sides of (2.1) to get

$$\sum_{k=1}^{\infty} \frac{H_{k-1}^{(2)} x^{k-1}}{k \binom{2k}{k}} = \frac{8}{3} \arcsin^3 \frac{\sqrt{x}}{2} \times \frac{1/(4\sqrt{x})}{\sqrt{1-x/4}} \quad \text{with } 0 \leq x < 4.$$

So (1.4) holds. Applying (1.4) with $x = 1, 3$ we obtain (1.5) and (1.6) respectively.

The proof of Theorem 1.1 is now complete. \square

Proof of Theorem 1.2. Let us recall a known hypergeometric identity

$$\sin(az) = a \sin(z) \cdot {}_2F_1 \left(\frac{1+a}{2}, \frac{1-a}{2}, \frac{3}{2}; \sin^2 z \right).$$

(See, e.g., [BE, §2.8].) For $k = 1, 2, 3, \dots$, clearly

$$\begin{aligned} & \frac{\prod_{j=0}^{k-1} ((1+a)/2 + j)((1-a)/2 + j)}{k! \prod_{j=1}^k (1/2 + j)} \\ &= \frac{\prod_{j=0}^{k-1} ((2j+1)^2 - a^2)}{k! (2k+1)! 2^k} = \frac{\binom{2k}{k}}{(2k+1)4^k} \prod_{j=0}^{k-1} \left(1 - \frac{a^2}{(2j+1)^2} \right). \end{aligned}$$

Therefore

$$\sin(az) = a \sin(z) \left(1 + \sum_{k=1}^{\infty} \frac{\binom{2k}{k}}{2k+1} \left(\frac{\sin^2 z}{4} \right)^k \prod_{j=1}^k \left(1 - \frac{a^2}{(2j-1)^2} \right) \right).$$

Comparing the coefficients of a^3 on both sides we get

$$\frac{z^3}{6} = \sin(z) \sum_{k=1}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)}}{2k+1} \left(\frac{\sin^2 z}{4} \right)^k. \quad (2.2)$$

Now let $0 < x \leq 4$ and set $z = \arcsin(\sqrt{x}/2)$. Then (2.2) gives (1.7).

Applying (1.7) with $x = ((\sqrt{5} \pm 1)/2)^2$ we obtain

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)}}{(2k+1)16^k} \left(\frac{\sqrt{5} \pm 1}{2} \right)^{2k+1} = \frac{1}{3} \left(\frac{(2 \pm 1)\pi}{10} \right)^3$$

and hence (1.8) and (1.9) follow.

Now prove (1.10) for $x \geq 1$. Note that

$$\sin(iz) = \frac{e^{-z} - e^z}{2i} = i \sinh(z),$$

where $\sinh(z)$ is the hyperbolic sine function. Substituting iz for z in (2.2) we get

$$-\frac{z^3}{6} = \sinh(z) \sum_{k=1}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)}}{2k+1} \left(-\frac{\sinh^2(z)}{4} \right)^k. \quad (2.3)$$

Fix $x \geq 1$ and let

$$\begin{aligned} z = \operatorname{arcsinh} \frac{1}{\sqrt{x}} &= \log \left(\frac{1}{\sqrt{x}} + \sqrt{1 + \frac{1}{x}} \right) \\ &= \log \frac{\sqrt{x+1} + 1}{\sqrt{x}} = \log \frac{\sqrt{x}}{\sqrt{x+1} - 1} = -\log \frac{\sqrt{x+1} - 1}{\sqrt{x}}. \end{aligned}$$

Then (2.3) yields (1.10).

So far we have completed the proof of Theorem 1.2. \square

Proof of Corollary 1.2. By Theorem 1.2(i), for $0 \leq x \leq 2$ we have

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)} x^{2k+1}}{(2k+1)16^k} = \frac{1}{3} \operatorname{arcsin}^3 \frac{x}{2}. \quad (2.4)$$

Observe that $\binom{2k}{k} < (1+1)^{2k} = 4^k$ and

$$\bar{H}_k^{(2)} < \sum_{j=1}^{\infty} \frac{1}{(2j-1)^2} = \sum_{j=1}^{\infty} \frac{1}{j^2} - \sum_{j=1}^{\infty} \frac{1}{(2j)^2} = \frac{\pi^2}{8}$$

for every $k = 1, 2, 3, \dots$. Let $0 < \varepsilon \leq 2$. Clearly the series

$$S(\varepsilon) := \sum_{k=1}^{\infty} \frac{4^k (2-\varepsilon)^{2k+1}}{(2k+1)16^k}$$

converges, and

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k} \bar{H}_k^{(2)} x^{2k+1}}{(2k+1)16^k} < \frac{\pi^2}{8} S(\varepsilon) \quad \text{for } 0 \leq x \leq 2 - \varepsilon.$$

Thus, we can take the derivatives of both sides of (2.4) with $0 \leq x \leq 2 - \varepsilon$ to obtain (1.17). Therefore (1.17) holds for any $0 \leq x < 2$.

Applying (1.17) with $x = 1, \sqrt{2}, \sqrt{3}$ we get (1.18)-(1.20) immediately. This concludes the proof. \square

REFERENCES

- [AKP] G. Almkvist, C. Krattenthaler and J. Petersson, *Some new series for π* , Experiment. Math. **12** (2003), 441–456.
- [BBB] D. H. Bailey, J. M. Borwein and D. M. Bradley, *Experimental determination of Apéry-like identities for $\zeta(2n+2)$* , Experiment. Math. **15** (2006), 281–289.
- [BE] H. Bateman and A. Erdélyi, *Higher Transcendental Functions*, Vol. 1, McGraw-Hill, New York, 1953.

- [BB] J. M. Borwein and D. H. Bailey, *Mathematics by Experiment: Plausible Reasoning in the 21st Century*, 2nd Edition, A K Peters, Wellesley, 2003.
- [DK] A. I. Davydychev and M. Yu. Kalmykov, *Massive Feynman diagrams and inverse binomial sums*, Nucl. Phys. B **699** (2004), 3–64.
- [M] R. Matsumoto, *A collection of formulae for π* , on-line version is available from the website http://www.pluto.ai.kyutech.ac.jp/plt/matsumoto/pi_small.
- [PS] H. Pan and Z. W. Sun, *A combinatorial identity with application to Catalan numbers*, Discrete Math. **306** (2006), 1921–1940.
- [PP] K. H. Pilehrood and T. H. Pilehrood, *Series acceleration formulas for beta values*, Discrete Math. Theor. Comput. Sci. **12** (2010), 223–236.
- [PWZ] M. Petkovšek, H. S. Wilf and D. Zeilberger, *A = B*, A K Peters, Wellesley, 1996.
- [S1] Z. W. Sun, *Binomial coefficients, Catalan numbers and Lucas quotients*, Sci. China Math. **53** (2010), 2473–2488.
- [S2] Z. W. Sun, *On congruences related to central binomial coefficients*, preprint, arXiv:0911.2415. <http://arxiv.org/abs/0911.2415>.
- [S3] Z. W. Sun, *On sums of binomial coefficients modulo p^2* , preprint, arXiv:0910.5667. <http://arxiv.org/abs/0910.5667>.
- [S4] Z. W. Sun, *Super congruences and Euler numbers*, preprint, arXiv:1001.4453. <http://arxiv.org/abs/1001.4453>.
- [S5] Z. W. Sun, *Correspondence between series for π and congruences for Euler numbers*, A Message to Number Theory List (sent on March 15, 2010)., <http://listserv.nodak.edu/cgi-bin/wa.exe?A2=ind1003&L=nmbirthry&T=0&P=906>.
- [S6] Z. W. Sun, *A new series for π^3 and related congruences*, preprint, arXiv:1009.5375. <http://arxiv.org/abs/1009.5375>.
- [ST1] Z. W. Sun and R. Tauraso, *New congruences for central binomial coefficients*, Adv. in Appl. Math. **45** (2010), 125–148.
- [ST2] Z. W. Sun and R. Tauraso, *On some new congruences for binomial coefficients*, Int. J. Number Theory, to appear. <http://arxiv.org/abs/0709.1665>.
- [Z] D. Zeilberger, *Closed form (pun intended!)*, Contemporary Math. **143** (1993), 579–607.